


SENSAR® 1-Piece Hydrophobic Acrylic IOL









DESCRIPTION	
OPTIC CHARACTERISTICS	
Powers:	+6.0 D to +30.0 D in 0.5 diopter increments
Diameter:	6.0 mm
Material:	UV-blocking hydrophobic acrylic
Refractive Index:	1.47
Edge Design:	ProTEC frosted, continuous 360° posterior square edge
ULTRASOUND BIOMETRY*	
A-constant:	118.4
Theoretical AC Depth:	5.2 mm
Surgeon Factor:1	1.45 mm
HAPTIC CHARACTERISTICS	
Overall Length:	13.0 mm
Style:	C
Material:	UV-blocking hydrophobic acrylic
Design:	Haptics offset from optic
RECOMMENDED INSERTION INSTRUMENTS	MODEL
UNFOLDER® Platinum 1 Series Handpiece UNFOLDER® Platinum 1 Series Cartridge One Series Ultra Implantation System (Syringe-style Inserter) One Series Ultra Implantation System (Screw-style Inserter) One Series Ultra Cartridge	DK7796 1MTEC30 DK7786 DK7791 1VIPR30

*Value theoretically derived for a typical 20.0 D lens. AMO recommends that surgeons personalize their A-constant based on their surgical techniques and equipment, experience with the lens model, and postoperative results.

To learn more and to view important safety information, please contact your AMO Sales Representative.

AMO United Kingdom Limited • Jupiter House • Mercury Park • Wycombe Lane Wooburn Green • High Wycombe • Buckinghamshire • HP10 OHH • UK

SENSAR® 1-piece lenses are indicated for the visual correction of aphakia in adult patients in whom a cataractous lens has been removed by extracapsular cataract extraction. These devices are intended to be placed in the capsular bag. For a complete listing of precautions, warnings, and adverse events, refer to the package insert.

One Series, SENSAR®, ProTEC and UNFOLDER® are trademarks owned by or licensed to Abbott Laboratories, its subsidiaries or affiliates.

©2011 Abbott Medical Optics Inc., Santa Ana, CA 92705 www.AbbottMedicalOptics.com 2011.11.14-CT4255



^{1.} Calculated based on Holladay I formula (Holladay JT, Prager TC, Chandler TY, Musgrove KH, Lewis JW, Ruiz RS. A three-part system for refining intraocular lens power calculations. J Cataract Refract Surg. 1988;14(1):17-24).